

OS NOV 86

DRAFT--DO NOT QUOTE OR CITE

main disadvantages of the study are: (1) it took place over a relatively narrow time frame of 8.5 years which increases the potential for a calendar year bias, and (2) the lower social economic classes are under-represented which may introduce a bias due to the difference in type and way cigarettes are smoked.

D.3.2.1.3. British male physicians. Doll and Peto (1978) published the data shown in Table D-6 based on the information obtained by following the survival of a cohort of approximately 34,000 British male physicians. The smoking histories of each individual in the cohort were obtained by questionnaires at three different points in time. Table D-6 is a subset of the total cohort consisting of subjects who smoked at a nearly constant rate over their smoking lifetime. Due to the quality of the smoking information and pathology confirmation of most of the cases, this study is generally acknowledged to be the most informative available for establishing dose-response relationships. The disadvantages are that the number of observed lung cancer deaths are relatively small (i.e., 215), no women are included in the sample, and information on ex-smokers was never published in a form suitable for analysis. Also, a sample of physicians has a high potential for a sociological bias to be built into it. Ten years of additional observation is available on the cohort that has not yet been published and could be of considerable importance in the establishment of a dose-response model.

TABLE D-6 NUMBER OF LUNG CANCER DEATHS AND PERSON-YEARS OF OBSERVATION FOR  
BRITISH MALE PHYSICIANS

Mid point age interval (years)		Average exposure (cigarettes per day)								
		0.0	2.7	6.6	11.3	16.0	20.4	25.4	30.2	38.0
42.5	observed lung cancer deaths	0.0	0.0	0.0	1.0	0.0	1.0	0.0	1.0	0.0
	person-years	17,846.5	1,216.0	2,041.5	3,795.5	4,824.0	7,046.0	2,523.0	1,715.5	892.5
47.5	observed lung cancer deaths	0.0	0.0	0.0	1.0	1.0	1.0	2.0	2.0	0.0
	person-years	15,832.5	1,000.5	1,745.0	3,205.0	3,995.0	6,460.5	2,565.5	2,123.0	1,150.0
52.5	observed lung cancer deaths	0.0	0.0	0.0	2.0	4.0	6.0	3.0	3.0	3.0
	person-years	12,226.0	853.5	1,562.5	2,727.0	3,278.5	5,583.0	2,620.0	2,226.5	1,281.0
57.5	observed lung cancer deaths	2.0	1.0	0.0	1.0	0.0	8.0	5.0	6.0	4.0
	person-years	8,905.5	625.0	1,355.0	2,288.0	2,428.5	4,357.5	2,108.5	1,923.0	1,063.0
62.5	observed lung cancer deaths	0.0	1.0	1.0	1.0	2.0	13.0	4.0	11.0	7.0
	person-years	6,248.0	509.5	1,068.0	1,714.0	1,829.5	2,863.5	1,508.5	1,362.0	826.0
67.5	observed lung cancer deaths	0.0	0.0	1.0	2.0	2.0	12.0	5.0	9.0	9.0
	person-years	4,351.0	392.5	843.5	1,214.0	1,237.0	1,930.0	974.5	763.5	515.0
72.5	observed lung cancer deaths	1.0	1.0	2.0	4.0	4.0	10.0	7.0	2.0	5.0
	person-years	2,723.5	242.0	696.5	862.0	683.5	1,055.0	527.0	317.5	233.0
77.5	observed lung cancer deaths	2.0	0.0	0.0	4.0	5.0	7.0	4.0	2.0	2.0
	person-years	1,772.0	208.5	517.5	547.0	370.5	512.0	209.5	130.0	88.5

Source: Doll and Peto (1978)

DRAFT--DO NOT QUOTE OR CITE

06 NOV 90

D-17

05/17/90

0822406202

05 NOV 90

DRAFT--DO NOT QUOTE OR CITE

D.3.2.1.4. Other data sources. Other sources of information that could prove useful in obtaining information on a dose-response model are Best (1966), Canadian smokers; Bross et al. (1968), individuals who switched to filter cigarettes; Cederlof et al. (1975), a national probability sample of Swedish subjects; Graham and Levin (1971), individuals who stopped smoking; Hirayama (1977), Japanese smokers; Stevens and Moolgavkar (1984), British males; Lubin et al. (1984), individuals who changed smoking habits; Wald et al. (1988), U.K. smoking statistics; the IARC monograph on the evaluation of the carcinogenic risk of tobacco smoking to humans IARC (1986), general information; and the U.S. Public Health Service, Smoking and Health Report series for various types of smoking related information.

D.3.2.2. Modeling Approach for Cigarette Smoking Data. Various investigators, such as Doll and Peto (1978), Thorslund and Charnley (1987), Brown and Chu (1987), Gailfney and Altshuler (1988), Darby and Pike (1988), Freedman and Navidi (1989), and Moolgavkar et al. (1989), were successful in fitting various forms of multi-stage type models to the British physicians data. Modeling attempts using the AWW and ACS data have been less successful. Freedman and Navidi (1989) could not obtain adequate fits using standard multi-stage models to the AWW and ACS data sets when information on ex-smokers was included. The reasons for this inability could be either deficiencies in the multi-stage model (hypothesis put forth by the authors) or some unknown bias in the data that distorts the true dose-response relationship. To clarify the situation other modeling approaches should be attempted.

Probably the most successful approach for mathematically modeling cigarette smoking data was put forth by Moolgavkar et al. (1989). This is the only attempt to date to incorporate a promotional component of cigarette smoke into a dose-response model. Using Moolgavkar's basic model and the additional simplifying robust assumptions that:

4 06 NOV 90

DRAFT--DO NOT QUOTE OR CITE

- the number of stem (target) cells are constant over time,
- the ratio  $S$  of unit exposure induced to background cell transition rates are equal for the two cellular transitions (i.e., normal stem to preneoplastic and preneoplastic to neoplastic), and
- the growth rate of preneoplastic cells is a function,  $G(x)$ , of the number of cigarettes smoked per day,

the age-specific lung cancer rate of an individual at age  $t$  who has smoked  $x$  cigarettes per day since age  $t_0$  can be expressed as:

$$h(x,t) = \frac{A}{G(x)}(1+Sx)^2[\exp\{G(x)(t-t_0)\}-1] + \frac{A}{G(0)}(1+Sx)\exp\{G(x)(t-t_0)\}[\exp\{G(0)t_0\}-1]$$

where  $A$  is the product of the background transition rates,  $G(x) = G(0)[1 + (R-1)M(x)]$  with  $R = G(\infty)/G(0)$  being the maximum relative growth rate increase that can be induced by cigarettes and  $M(x)$  is a still to be specified function that defines the fraction of the maximum growth rate increase that is induced with  $x$  cigarettes per day. In the model employed by Moolgavkar, the simplifying assumption  $G(x) = G(0) + \Delta x$  was made. While this assumed relationship may be appropriate at low doses, it very likely results in a distortion of the effect for heavy smokers.

It is proposed that the Moolgavkar (i.e., two-stage) model parameter estimates be obtained by simultaneously using multiple epidemiological-smoking-lung cancer data sets and the following modifications and extensions of the above basic model:

- Moolgavkar assumed that the time from the development of a neoplastic cell until death due to a lung cancer was a constant 3.5 years for each of the lung cancer deaths. As an alternative this length of time will be estimated by maximum likelihood methods assuming:
  1. it is a constant unknown value for all lung cancer deaths, and
  2. it is a random variable with an integer gamma probability distribution.
- Alternative specific forms for  $G(x)$  will be specified based on various assumptions of how binding of smoking product agents with preneoplastic cells induce promotion.

05 NOV 90

DRAFT--DO NOT QUOTE OR CITE

- An adjustment will be made for the difference between British and American cigarettes and British and American smoking habits.
- An investigation will be made of the hypothesis that  $G(0)$  may be different pre- and post-exposure to accommodate the observation of rapidly falling age-specific rates post cessation of smoking.

The largest information data base possible will be used in fitting the different variations of the model. An illustration of how one of the parameters in the model,  $G(0)$ , could be estimated is given below.

Hammond (1966) pooled the ACS lung cancer mortality data for men and women nonsmokers and obtained age-specific death rates for five-year age intervals. This information is displayed in Table D-7. The justification given by Hammond (1966) for pooling the data was the inability to reject the hypothesis of equal rates for the sexes on the basis of a statistical test. Under the assumption of no cigarette smoking,  $x=0$ , so the previously described age-specific rates for the two-stage model has the reduced form:

$$M(0,A) = \frac{A}{G(0)} [\exp(G(0)A) - 1]$$

Assuming that the number of lung cancer cases out of the number of person years of observation was an independent binomial random variable for each age class, maximum likelihood estimates were obtained for the unknown parameters  $A$  and  $G(0)$  in the above model. The adequate fit of the model is displayed in Table D-7 and Figure D-1.

It is reasonable to assume that the parameter  $G(0)$  is human population independent and, perhaps, even species independent taken on a lifetime equivalent time scale. However, the value  $A$  would most likely be dependent on the environmental conditions an individual is living under. Therefore, different values for U.S. and British populations should be estimated.

TABLE D-7. LUNG CANCER DEATH RATES PER 100,000 PERSON-YEARS AND  
OBSERVED AND PREDICTED NUMBER OF LUNG CANCER DEATHS AMONG  
MEN AND WOMEN WHO NEVER SMOKED REGULARLY

Age group L to L+5	Combined men and women			Population size N (person-years)
	Number of lung cancer deaths n		Death rate dr	
	Observed	Predicted		
40-44	4	5.40	2.3	173,913
45-49	16	14.01	5.0	320,000
50-54	16	20.05	4.9	326,531
55-59	30	24.52	10.5	285,714
60-64	32	27.53	13.9	230,216
65-69	26	29.43	14.7	176,871
70-74	18	25.84	16.1	111,801
75-79	21	18.82	35.8	58,659
80-84	14	11.41	54.6	25,641
Total	177	----		

Source: Hammond (1966)/ACS Study

$$N = \frac{n \times 10^6}{dr}$$

calculated from data  $\chi^2 = 7.036$   $p = 0.425$

DRAFT--DO NOT QUOTE OR CITE

D-21


05/17/90

4822406202

4 05 NOV 80

DRAFT--DO NOT QUOTE OR CITE

FIGURE D-1.  
GOODNESS-OF-FIT OF TWO-STAGE MODEL TO NON-SMOKERS  
AGE-DEPENDENT LUNG CANCER DATA



Source of data: Hammond et al. (1966).

The obvious advantage of the proposed CEA is that it is based on the most extensive body of information concerning the dose-dependent effects of an environmental agent on a human cancer response that exists. The main disadvantages are the complexity of the analysis and the possibility of not establishing a credible ETS equivalency relationship. The latter factor is discussed in the next sections.

#### D.3.3. Estimation of the Relative Potency of ETS Compared to MS

Previous approaches for establishing ETS/cigarette equivalency (e.g., Darby and Pike, 1988) have made the implicit assumption that the ratio of the potency of emissions to some surrogate measure of internal exposure (e.g., nicotine, cotinine, etc.) is the same for ETS and MS. The large variability in relative potency estimates of complex-PAH mixtures that are displayed in Table D-8 suggests that the implicit assumption of equal potency is suspect.

Several methods can be used to estimate the ETS compared to MS relative potency. The inhalation studies in Syrian golden hamsters where laryngeal carcinomas were elicited from MS (Dontenwill et al., 1973; 1977) and from B[a]P (Thyssen, 1981) can be used to obtain a MS-to-B[a]P relative potency estimate. Dividing this obtained potency value into the ETS-to-B[a]P, the relative potency obtained from the lung implant studies discussed in Section D.1.3.1 would give a relatively potent estimate of ETS to MS. Stanton et al. (1972) conducted a lung implant study using cigarette smoke condensate (CSC). Unfortunately for our present purposes, 3-methylcholanthrene (MCA) was used as the positive control in the experiment so direct comparison with ETS is not possible. However, Grimmer and his colleagues for the most part closely adopted Stanton's experimental protocol for conducting lung implant studies. Thus, a direct pooling of the data in the Stanton and Grimmer experiments could logically be used to obtain a potency estimate. As an alternative, the two-step approach of estimating the potency of



4 06 NOV 90

DRAFT--DO NOT QUOTE OR CITE

TABLE D-8. RELATIVE POTENCY ESTIMATES OF COMPLEX MIXTURES  
OF INCOMPLETE COMBUSTION PRODUCTS OF HYDROCARBONS  
COMPARED TO B[a]P

Complex PAH exposure	Direct bioassay estimate* of relative potency
Coal Flue Gas Condensate	0.05444
Gas Engine Condensate	0.02190
Diesel Engine Exhaust	0.00230
Sidestream Cigarette Smoke	0.00302
Coke Oven Emissions	0.03180 <sup>†</sup>

\*Lung implant studies

<sup>†</sup>Skin painting

CSC compared to MCA from the Stanton experiment and then establishing the relative potency of MCA compared to B[a]P in another assay system (e.g., subcutaneous injection, skin painting, etc.) could be employed. A final alternative might be to compare the weighted relative potency estimates of the known constituents in the MS and ETS samples that have stable established estimates of their carcinogenic potency compared to B[a]P. One potential list of stable relative potency estimates developed by Thorslund (1990) is shown in Table D-9.

The last piece of information required to obtain an ETS risk model based upon the CEA is a deposition ratio estimate between MS via active smoking and ETS under normal inhalation conditions. One promising approach of using B[a]P-DNA-adducts and other endpoints as biomarkers is discussed in the next section.

#### D.3.4. Deposition Differences of Chemicals from Cigarette Smoke in Smokers and Nonsmokers

To obtain an equivalency relationship between MS and ETS, both potency and deliverable dose conversion factors are needed in order to use the MS-lung cancer data as a surrogate for lung cancer induced by ETS.

06 NOV 90

DRAFT--DO NOT QUOTE OR CITE

TABLE D-9. RELATIVE POTENCY ESTIMATES OF AGENTS  
COMPARED TO B[a]P

Agent	Relative potency	Source of estimate
Anthracene	0.00000	IARC adequately studied, no indication of carcinogenic effect category
Fluoranthene	0.00000	
Pyrene	0.00000	
Benzo[b]fluoranthene	0.12277	Deutsch-Wenzel et al. (1983) (Grimmer's group) lung implant data Thorslund (1990) estimates
Benzo[k]fluoranthene	0.05322	
Benzo[j]fluoranthene	0.05232	
Benzo[e]pyrene	0.00704	
Benzo[a]pyrene	1.00000	
Indeno Pyrene	0.27800	
Benzo[ghi]perylene	0.02124	
Anthanthrene	0.31598	

Under the assumption that the PAHs possess most of the carcinogenic potency in MS and ETS, the deliverable target dose can be estimated by directly measuring the number of DNA adducts formed in people smoking different numbers of cigarettes per day and in people who are nonsmokers in the presence of smokers with different frequencies of smoking.

Specific adducts, such as the DNA 7,8-diol-9,10-epoxide of B[a]P which is present in both MS and ETS, can be detected using sensitive immunoassays or postlabelling DNA techniques (Shamsuddin et al., 1985; Randerath et al., 1986). Differences in adduct formation between smokers and nonsmokers varied depending on the experiment but was as high as 400-fold when DNA from oral mucosa was analyzed using the postlabelling technique. Hemoglobin adducts as markers of genotoxicity have been analyzed in smokers and nonsmokers where smokers had about a 7-fold greater number of adducts than nonsmokers.

05 NOV 80

DRAFT--DO NOT QUOTE OR CITE

Indirect measures of dose between smokers and nonsmokers may also be available in which gene mutations can be measured in peripheral leukocytes at the Hypoxanthine phosphoribosyl transferase locus as well as other loci. In fact, such a test could conceivably be used directly to obtain a cigarette equivalence estimate without making potency difference adjustments. Other genetic damage tests, such as chromosomal aberrations and sister chromatid exchanges, may also be useful in determining deliverable target dose information for smokers and nonsmokers exposed to ETS.

To obtain an equivalency relationship of deliverable dose between smokers and nonsmokers, a thorough review of the literature for articles that show dose-response relationships between MS/ETS and DNA adducts, protein adducts, and gene mutations should be conducted and the most appropriate endpoints selected for use in the equivalency estimate. The main advantage of the approach is the high suspected correlation of the endpoint with the cancer response. The main disadvantage is the discounting of potential agents that act exclusively as promoters.

#### D.4. DIRECT APPROACH

The most straightforward approach for estimating ETS lung cancer risk is to estimate ETS exposure in a suitable cohort and follow the resulting mortality pattern over time. As of yet, no directly measured ETS exposure data exist on a cohort. The ideal in this regard would be personal monitoring data obtained from nonsmokers for an agent such as cotinine which is closely and uniquely associated with cigarette smoke. In this application, the use of cotinine is appropriate as long as it is linearly related to total ETS air levels. In lieu of such information, investigators have attempted to obtain surrogate measures of ETS. One such measure is the number of cigarettes smoked per day by the spouses of nonsmoking individuals. The quality of such a surrogate measurement depends upon: (1) the extent that nonsmokers are exposed to

DRAFT--DO NOT QUOTE OR CITE

smokers other than their spouses, (2) the consistency within the cohort of the husbands' and wives' spatial and time closeness, and (3) the consistency within the cohort of the fraction of the total cigarettes that are smoked by the spouse in the home. Due to sociological factors regarding a woman's place in Japan, the homogeneity of the Japanese society, and the small, close living arrangements of Japanese couples, probably the best surrogate measure of ETS exposure available is the number of cigarettes smoked per day by the husbands of Japanese women. The person-years of observation and the number of lung cancer deaths for Japanese women classified in regard to their husband's age and smoking habits obtained in the prospective study conducted by Hirayama (1984) is displayed in Table D-10. Under the assumption that all the excess lung cancer risk in Japanese women was due to husband-produced ETS exposure in the home, crude risk models can be generated from the information supplied in Table D-10. Better estimates could be obtained if information such as the length of marriage, wife's age, age husband started smoking, and smoking habits of wife's parents were available for individual cohort members. A fair amount of such information has been generated by Hirayama (1984) but presently is not reported in the open literature. Gaining access to the data could prove valuable.

17 05 NOV 50

TABLE D-10. LUNG CANCER MORTALITY IN JAPANESE WOMEN BY HUSBAND'S AGE GROUP AND SMOKING HABITS (PATIENT HERSELF A NON-SMOKER)\*

Husband's age group	Husband's smoking habit											
	Non-smoker		Ex-smoker		1-14/day		15-19/day		20+ /day		Total	
40-49	4	6,229	1	1,255	8	8,621	6	5,158	16	10,764	35	32,027
50-59	10	7,791	3	1,922	20	9,668	8	4,052	24	9,820	65	33,253
60-69	15	7,120	11	2,687	28	7,243	9	2,513	23	4,651	89	24,214
70-79	5	755	2	348	2	612	1	105	1	226	11	2,046
Total	37	21,895	17	6,212	58	26,144	24	11,828	64	25,461	200	91,540

\*Number of lung cancer deaths out of number of wives in the same cross classification cell.  
Source: Hirayama (1984).

DRAFT-DO NOT QUOTE OR CITE

D-28

05/17/91

1622406202

06 NOV 90

DRAFT--DO NOT QUOTE OR CITE

#### REFERENCES AND RELATED BIBLIOGRAPHY

- Agresti, A. (1990) *Categorical data analysis*. John Wiley & Sons, Inc.
- Akiba, S.; Kato, H.; Blot, W.J. (1986) Passive smoking and lung cancer among Japanese women. *Cancer Research* 46:4804-4807.
- Albert, R.E.; Lippman, M.; Briscoe, W. (1969) The Characteristics of bronchial clearance in humans and the effect of cigarette smoking. *Arch. Environ. Health* 18:738-755.
- Albert, R.E.; Peterson, H.T.; Bohning, D.E.; Lippmann, M. (1975) Short-term effects of cigarette smoking on bronchial clearance in humans. *Arch. Environ. Health* 30:361-367
- Albert, R.E., Lewtas, J., Nesnow, S., Thorslund, T., Anderson, E. (1982) Comparative potency method for cancer risk assessment: Application to diesel particulate emissions. *J. Risk Anal* 101-107.
- Altschuler, B.; Nelson, N.; Kuschner, M. (1964) Estimation of lung tissue dose from the inhalation of radon and daughters. *Health Phys.* 10:1137-1161.
- American Academy of Pediatrics, Committee on Environmental Hazards (1986) Involuntary smoking--A hazard to children. *Pediatrics* 77(5):755.
- Angioni, A.M.; Fanciulli, G.; Corchia, C. (1989) Frequency of and risk factors for allergy in primary school children: Results of a population survey. *Pediatric and Perinatal Epidemiology* 3:248-255.
- Air Pollution Control Association (APCA) (1986) *Indoor Radon*, APCA, Pittsburgh, PA.
- Aronow, W.S. (1978) Effect of passive smoking on angina pectoris. *New England Journal of Medicine* 299(1):21-24.
- Arundel, A.; Sterling, T.; Weinkam, J. (1987) Never smoker lung cancer risks from exposure to particulate tobacco smoke. *Environ. Int.* 13:409-426.
- Badre, R.; Guillermin, R.; Abran, N.; Bourdin, M.; Duman, C. (1978) Atmospheric pollution by cigarette smoking. *Ann Pharm Fr.* 36:443-452.
- Balter, N.J.; Schwartz, S.L.; Kilpatrick, S.J.; Witorsch, P. (1986) Causal relationship between environmental tobacco smoke and lung cancer in non-smokers: a critical review of the literature. *Proceedings of the Air Pollution Control Association, 79th annual meeting*.
- Banks, M.H.; Bewley, B.R.; Bland, M.; Pollard, V. (1978) Effect of children's and parents' smoking on respiratory symptoms. *Archives of Disease in Childhood* 53(2) 100-105.
- Barker, D.; Osmond, C. (1986) Childhood respiratory infection and adult chronic bronchitis in England and Wales (UK). *British Medical Journal* 293(6557):1271-1275.

05 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- Bassi, J.A.; Rosso, P.; Moessinger, A.C.; Blanc, W.A.; James, L.S. (1984) Fetal growth retardation due to maternal tobacco smoke exposure in the rat. *Pediatric Research* 18(2):127-130.
- Battista, S.P.; Kensler, C.J., (1970) "Mucus Production and Ciliary Transport Activity." *Arch. Environ. Health* 20:326-338.
- Benowitz, M.L.; Jacob, P.; Jones, R.T.; Rosenberg, J. (1982) Inter-individual variability in the metabolism and cardiovascular effects of nicotine in man. *J. Public Health* 221:368-372.
- Benowitz, N.L.; Jacob, P. (1984) Daily intake of nicotine during cigarette smoking. *Clin. Pharmacol. Ther.* 35:499-504.
- Bergman, H.; Axelson, A. (1983) Passive smoking and indoor radon daughter concentration. *Lancet* 8362:1308-1309.
- Bergman, H.; Edling, C.; Axelson, O. (1986) Indoor Radon Concentrations and Passive Smoking. *Environ. Int.* 12:17-19.
- ✓ Berkey, C.S.; Dockery, D.W.; Ferris, B.G.Jr.; Speizer, F.E.; Ware, J.H. (1986) Indoor air pollution and pulmonary function growth in preadolescent children. *American Journal of Epidemiology* 123(2):250-260.
- Berkey, C.S.; Ware, J.H.; Speizer, F.E.; Ferris, B.G.Jr. (1984) Passive smoking and height growth of preadolescent children. *International Journal of Epidemiology* 13(4):454-458.
- Best, E.W.R. (1966) A Canadian study of smoking and health. Department of National Health and Welfare, Ottawa.
- Bewley, B.R.; Bland, J.M.; Harris, R. (1974) Factors associated with the starting of cigarette smoking by primary children. *British Journal of Preventive Social Medicine* 28:37-44.
- Black, N. (1978) The aetiology of glue ear: A case-control study. *International Journal of Pediatric Otorhinolaryngology* 9(2):121-133.
- Black A.; Pritchard, J. N. (1984) A comparison of the regional deposition and short-term clearance of tar particulate material from cigarette smoke, with that of 2.5  $\mu$ m polystyrene spheres. *Aerosol Sci.* 15:224-227.
- Bland, M.; Bewley, B.R.; Pollard, V.; Banks, M.H. (1978) Effect of children's and parents' smoking on respiratory symptoms. *Archives of Disease in Childhood* 53(2):100-105.
- Blot, W.J.; Fraumeni, J.F. (1986) Passive smoking and lung cancer. *Journal of the National Cancer Institute* 77(5):993-999.
- Blot, W.J.; McLaughlin, J.K. (1985) Practical issues in the design and conduct of case-control studies: use of next-of-kin interviews. Blot, W.J., Hirayama, T., Huel, O.G., eds *Statistical Issues in Cancer Epidemiology*. Hiroshima: Sanei, pp. 46-62.

05 NOV 90

DRAFT--DO NOT QUOTE OR CITE

- Breslow, N.E.; Day, N.E. (1980) Statistical methods in cancer research. Lyon: IARC Scientific Publications No. 32.
- Bross, I.D.J.; Gibson, R. (1968) Risks of lung cancer in smokers who switch to filter cigarettes. *Am J Publ Hlth* 58:1369-1403.
- Brown, C.C.; Chu, K. (1987) Use of multistage models to infer stage affected by carcinogenic exposure: Example of lung cancer and cigarette smoking. *J Chronic Dis* 40(Suppl):171S-179S.
- Brown, C.C.; Chu, K. (1987) Use of multistage models to infer stage affected by carcinogenic exposure: Example of lung cancer and cigarette smoking. *J. Chronic Dis.* 40(2):171A-179A.
- Brownson, R.C.; Reif, J.S.; Keefe, T.J.; Ferguson, S.W.; Pritzl, J.A. (1987) Risk factors for adenocarcinoma of the lung. *Am J Epidemiol* 125:25-34.
- Brunekreef, B.; Boelrij, J.S. (1982) Long-term average suspended particulate concentrations in smokers' homes. *Int Arch Occup Environ Health* 50:299-302.
- Brunekreef, B.; Fischer, P.; Remijn, B.; Van der Lende, R.; Schouten, J.; Quanjer, Ph. (1985) Indoor air pollution and its effects on pulmonary function of adult non-smoking women: III. Passive smoking and pulmonary function. *International Journal of Epidemiology* 14(2):227-230.
- Brunnemann, K.D.; Adams, J.D.; Ho, D.P.; Hoffmann D. (1978) The influence of tobacco smoke on indoor atmospheres. II. Volatile and tobacco-specific nitrosamines in main and sidestream smoke and their contribution to indoor pollution. *Proceedings of the 4th Joint Conference on Sensing of Environmental Pollutants*. Washington, DC. American Chemical Society, pp. 876-880.
- Brunnemann, K.D.; Hoffmann D. (1978) "Chemical studies on tobacco smoke LIX. Analysis of volatile nitrosamines in tobacco smoke and polluted indoor environments." Walker, E.A., Castegnaro, M., Gričute, L. and Lyle, R.E., eds. *Environmental aspects of N-nitroso compounds* Lyon: IARC Scientific Publications No. 19, pp. 343-356.
- Buffler, P.A.; Pickle, L.W.; Mason, T.J.; and Contant, C. (1984) The causes of lung cancer in Texas. Mizell, M. and Correa, P., eds. *Lung Cancer: Causes and Prevention*. New York: Verlag Chemie International, pp. 83-99.
- Burchfiel, C.M.; Higgins, M.W.; Keller, M.W.; Howatt, W.F.; Butler, W.J.; Higgins, I.T.T. (1986) Passive smoking in childhood - Respiratory conditions and pulmonary function in Tecumseh, Michigan. *Am. Rev. Respir. Dis.* 133:966-973.
- Cameron, P.; Kostin, J.S.; Oselett, B.; Stocker, R.; Tighe, G.; Winton, J.; Wolfe, J.H.; Zaks, J.M. (1969) The health of smokers' and nonsmokers' children. *Journal of Allergy* 43(6):336-341.



06 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- Camner, P.; Philipson, K.; Arvidsson, T. (1971) Cigarette smoking in man, short-term effect on mucociliary transport. *Arch. Environ. Health* 23:421-426.
- Canadian Pediatric Society (1986) Secondhand cigarette smoke worsens symptoms in children with asthma. *CMAJ* 135(4):321-323.
- Cano, J. P.; Catalin, J.; Bnadre, R.; Dumas, C.; Viala, A.; Guillermin, R. (1970) Analysis of nicotine by gas chromatography: II applications. *Ann Pharm Fr* 28:633-640.
- Carter, S.L.; Hasegawa. (1975) Fixation of tobacco smoke aerosols for size distribution studies. *J Colloid Interface Science* 53:134-141.
- Cedarlof, R., Friberg, L., Hurbec, Z., Lorch, U. (1975) The relationship of smoking and some social covariables in mortality and cancer morbidity: A ten year follow-up on a probability sample of 55,000 Swedish subjects age 18-69, Parts 1 and 2. Stockholm: Department of Environmental Hygiene, The Karolinska Institute.
- Chan, W.C.; Colbourne, M.J.; Fung, S.C.; Ho, H.C. (1979) Bronchial cancer in Hong Kong, 1976-1977. *Br. J. Cancer* 39:182.
- Chan, W.C.; Fung, S.C. (1982) Lung cancer in non-smokers in Hong Kong. *Cancer Campaign*, Vol 6, *Cancer Epidemiology*, (Grundmann, E., ed.) Stuttgart: Gustav Fischer Verlag, pp. 199-202.
- Chan, K.N.; Elliman, A.; Bruan, E.; Silverman, M. (1989) Respiratory symptoms in children of low birth weight. *Arch. Dis. Child.* 64(9):1294-1304.
- Chappell, S.B.; Pasker, R.J. (1977) Smoking and carbon monoxide levels in enclosed public places in New Brunswick. *Can J of Public Health* 68:159-161.
- Charlton, A. (1984) Children's coughs related to parental smoking. *British Medical Journal* 288:1647-1649.
- Charlton, A.; Blair, V. (1989) Absence from school related to children's and parental smoking habits. *British Medical Journal* 298:90-92.
- Checkoway, H.; Pearce, N.; Crawford-Brown, D. (1989) *Research methods in epidemiology*. New York, NY: Oxford University Press.
- Chen, Y.; Li, W.-X. (1986a) The effect of passive smoking on children's pulmonary function in Shanghai. *American Journal of Public Health* 76(5):515-518.
- Chen, Y.; Li, W.-X.; Yu, S. (1986b) Influence of passive smoking on admissions for respiratory illness in early childhood. *British Medical Journal* 293:303-306.
- Chen, Y.; Li, W.-X.; Yu, S.; Qian, W. (1988) Chang-Ning epidemiological study of children's health: I: Passive smoking and children's respiratory diseases. *International Journal of Epidemiology* 17(2):348-355.

4 06 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- Chen, Y. (1989) Synergistic effect of passive smoking and artificial feeding on hospitalization for respiratory illness in early childhood. *Chest* 95(5):1004-1007.
- Coburn, R.F.; Forster, R.E.; Kane, P.B. (1965) Considerations of the physiological variables that determine the blood carboxyhemoglobin concentration in man. *J Clin Invest* 44:1899-1910.
- Colley, J.R.T. (1971) Respiratory disease in childhood. *British Medical Bulletin* 27(1):9-14.
- Colley, J.R.T. (1974) Respiratory symptoms in children and parental smoking and phlegm production. *British Medical Journal* 2(5912):201-204.
- Comroe, J.H.; Forster II, R.E.; Dubois, A.B.; Briscoe, W.A.; Carlsen, E. (1962) The lung: Clinical physiology and pulmonary function tests. Chicago: Year Book Medical Publ. Inc., pp. 323-364.
- Comstock, G.W.; Helsing, K.J.; Meyer, M.B.; Tockman, M.S. (1981) Respiratory effects of household exposures to tobacco smoke and gas cooking. *American Review of Respiratory Disease* 124(2):143-148.
- Copeland, K.T.; Checkoway, H.; McMichael, A.J.; Holbrook, R.H. (1977) Bias due to misclassification in the estimation of relative risk. *Am J of Epidemiol.* 105(5):488-95.
- Cornfield, J. (1951) A method of estimating comparative rates from clinical data: Applications to cancer of the lung, breast, and cervix. *J. Nat'l Cancer Inst.* 11:1269-1275.
- Correa, P.; Fontham, E.; Pickle, L.; Lin, Y.; Haenszel, W. (1983) Passive smoking and lung cancer. *Lancet* 2:595-597.
- Correa, P. (1986) The passive smoking-cancer controversy. In: *Mechanisms in Tobacco Carcinogenesis*. Banbury Report 23:343-360.
- Cothern, C. (1987) "Properties," in *Environmental Radon*, edited by C. Cothern and J. Smith, Plenum Press, New York, p.1-29.
- Cothern, C.; Smith, J., editors, (1987) *Environmental Radon*, Plenum Press, New York.
- Coultas, D.B.; Howard, C.A.; Peake, G.T.; Skipper, B.J.; Samet, J.M. (1987) Salivary cotinine levels and involuntary tobacco smoke exposure in children and adults in New Mexico. *Am Rev Respir Dis* 136:305-309.
- Covey, L.S., Mushinski, M.H., Wynder E.L. (1983) Smoking habits in a hospitalized population: 1970-1980. *American Journal of Public Health* 73:1293-1297.
- Crawford, D.J. (1982) Identifying critical human subpopulations by age groups: radioactivity and the lung. *Phys. Med. Biol.* 27:539-552.
- Crawford-Brown, D.J. (1984) A unified approach to age-dependent metabolic modeling. *Health Physics* 46:809.

06 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- Crawford-Brown, D.J. (1987) Dosimetry. Cothorn C.R. and Smith, J.E., Jr., eds. Environmental Radon. New York, NY: Plenum Press, pp. 173-213.
- Crawford, D.J. (1982). Identifying critical human subpopulations by age groups: Radioactivity and the lung. *Phys. Med. Biol.* 27, 539-552.
- Crawford, D.J.; Eckerman, K. (1983) Modifications of the ICRD task group lung model to reflect age dependence. *Rad Prot Dosim* 2:209-220.
- Cuddeback, J.E.; Donovan, J.R.; Buring, W.R. (1976) Occupational aspects of passive smoking. *Am Ind Hyg Assoc* 37:263-267.
- Cummings, K.M.; Markello, S.J.; Mahoney, M.; Bhargava, A.K.; McElroy, P.D.; Marshall, J.R. (1986) Measurement of current exposure to environmental tobacco smoke. *Am J Epidemiol* 130(1):122-132.
- Cummings, K.M.; Markello, S.J.; Mahoney, M.C.; Marshall, J.R. (1989a) Measurement of lifetime exposure to passive smoke. *Am J Epidemiol* 30:122.
- Dahms, T.E.; Bolin, J.F.; Slavin, R.G. (1981) Passive smoking: Effects on bronchial asthma. *Chest* 80(5):530-534.
- Dalager, N.A.; Pickle, L.W.; Mason, T.J.; Correa, P.; Fontham, E.; Stemhagen, A.; Buffler, P.A.; Ziegler, R.G.; Fraumeni, J.F. (1986) The relation of passive smoking to lung cancer. *Cancer Research* 46:4808-4811.
- Darby, S.C.; Pike, M.C. (1988) Lung cancer and passive smoking: Predicted effects from a mathematical model for cigarette smoking and lung cancer. *Br J Cancer* 58:825-831.
- Deutsch-Wenzel, R.P.; Brune, H.; Grimmer, G.; Dettbarn, G.; Misfeld, J. (1983) Experimental studies in rat lungs on the carcinogenicity and dose-response relationships of eight frequently occurring environmental polycyclic aromatic hydrocarbons. *J Natl Cancer Inst* 71:539-544.
- Dockery, D.W.; Spengler, J.D. (1981) Indoor-outdoor relationships of respirable sulfates and particles. *Atmos Environ* 15:335-344.
- Dockery, D.W.; Spengler, J.D. (1981) Personal exposure to respirable particles and sulfates. *J. Air Pollut Control Assoc* 31:153-159.
- Dodge, R. (1982) The effects of indoor pollution on Arizona children. *Archives of Environmental Health* 37(3):151-155.
- Doll, R.; Peto, R. (1976) Mortality in relation to smoking: 20 years' observations on male British doctors. *Br. Med. J.* ii, 1525-1536.
- Doll, R.; Peto, R. (1978) Cigarette smoking and bronchial carcinoma: dose and time relationships among regular smokers and lifelong non-smokers. *J. Epidemiol. Commun. Health* 32: 303-313.

05 NOV 90

DRAFT--DO NOT QUOTE OR CITE

- Doll, R.; Peto, J. (1986) Passive smoking. *Br J Cancer* 54:1019-1021.
- Dontenwill, W.; Chevalier H.-J.; Harke H.-P.; Lafrenz, U.; Reckzeh, G.; Schneider, B. (1973) Investigations on the effects of chronic cigarette smoke inhalation in Syrian goldenhamsters. *J. Natl Cancer Inst* 51:1781-1832.
- Dontenwill, W.; Chevalier, H.-J.; Harke, H.-P.; Klimisch, H.-J.; Kuhnigle, C.; Reckzeh, G.; Schneider, B. (1977) Studies on the effect of chronic cigarette smoke inhalation in Syrian golden hamsters and the importance of vitamin A on morphological alterations after smoke exposure. *Z. Krebsforsch* 89:145-151.
- Ebert, R.V.; Terracio, M.J., (1975), The Bronchiolar Epithelium in Cigarette Smokers, *American Rev. Resp. Dis.* 111:4-11.
- Ekwo, E.E.; Huntley, W.H.; Lachenbruch, P.A.; Weinberger, M.M. (1983) Relationship of parental smoking and gas cooking to respiratory disease in children. *Chest* 84(6):662-668.
- Elliot, L.P.; Rowe, D.R. (1975) Air quality during public gatherings. *J. Air Pollut Control Assoc* 25:635-636.
- Eriksen, M.P.; LeMaistre, C.A.; Newell, G.R. (1988) Health hazards of passive smoking. *Ann Review of Public Health* 9:47-70.
- Eudy, L.W.; Thomas, F.A.; Heavner, D.L.; Green, C.R.; Ingebrethsen, B.J. (June 22-27, 1986) Studies on the vapor-particulate phase distribution of environmental nicotine by selective trapping and detection methods. *Proc of the Air Pollution Control Association 79th annual meeting*, pp. 2-14.
- Evans, R.E. (1968) Engineers guide to the elementary behavior of radon daughters. *Health Phys.* 38:1173-1197.
- Evans, D.; Levison, M.J.; Feldman, C.H.; Clark, N.M.; Wasilewski, D.; Levin, B.; Mellins, R.B. (1987) The impact of passive smoking on emergency room visits of urban children with asthma. *American Review of Respiratory Disease* 135:567-572.
- Federal Register (1986) Guidelines for carcinogen risk assessment. Vol. 51, No. 185, p. 33994-34003, Sept. 24, 1986. Also available as U.S. Environmental Protection Agency (EPA). (1987) The risk assessment guidelines of 1986. Office of Health and Environmental Assessment, Washington, DC. EPA/600/8-87/045.
- Fergusson, D.M.; Horwood, L.J.; Shannon, F.T.; Taylor, B. (1981) Parental smoking and lower respiratory illness in the first three years of life. *Epidemiology and Community Health* 35(3):180-184.
- Fergusson, D.M.; Hons, B.A.; Horwood, L.J. (1985) Parental smoking and respiratory illness during early childhood: A six-year longitudinal study. *Pediatric pulmonology* 1:99-106.
- Ferris, B.G.; Ware, J.H.; Berkey, C.S.; Dockery, D.W.; Spiro, A. III; Speizer, F.E. (1985) Effects of passive smoking on health of children. *Environmental Health Perspectives* 62:289-295.

06 NOV 80

DRAFT-DO NOT QUOTE OR CITE

- Gastineau, R.M.; Walsh, P.J.; Underwood, N. (1972) Thickness of bronchial epithelium with relation to exposure to radon. *Health Phys.* 23:857-860.
- Geller-Bernstein; Kenett, R.; Weisglas, L.; Tsur, S.; Lahav, M.; Levin, S. (1987) Atopic babies with wheezy bronchitis. *Allergy* 42:85-91.
- Geng, G.; Liang, Z.H.; Zhang, G.L. (1988) On the relationship between smoking and female lung cancer. *Smoking and Health*, Elsevier Science Publishers, pp. 483-86.
- George, A.; Knutson, E.; Tu, K.W. (1983) Radon Daughter Plateout. *Health Phys.* 45:439-444.
- Gies, R.A.; Cross, F.T.; Dagle, G.E. (1972) A histological study of the influence of cigarette smoking in suppressing Rn daughter carcinogenesis in dogs. *Health Phys.* 23:857-860.
- Gillis, C.R.; Hufe, D.J.; Hawthorne, V.M.; and Boyle, P. (1984) The effect of environmental tobacco smoke in two urban communities in the west of Scotland. *Eur J Respir Dis* 133(Suppl.):121-126.
- Godin, G.; Wright, G.; Shephard, R.J. (1972) Urban exposure to carbon monoxide. *Arch Environ Health* 25:305-313.
- Göldstein, G.M.; Collier, A.; Etzel, R.; Lewtas, J.; Haley, N. (1987) Elimination of urinary cotinine in children exposed to known levels of side-stream cigarette smoke. Seifert, H.; Esdorn, M.; Fischer, H.; Ruden, J.; Wegner, eds. In: *Indoor Air '87*, Vol 2, B. Oranienbruck GmbH, Berlin. pp. 61-67.
- Goodman, R. M.; Yergin, B. M.; Landa, J. F.; Golinvaux, M. H.; Sackner, M. A. (1978) Relationship of smoking history and pulmonary function tests to tracheal mucus velocity in non-smokers, young smokers, ex-smokers and patients with chronic bronchitis. *Am Rev Resp Dis* 117:205-214.
- Goodman, D.G.; Ward, J.M.; Squire, R.A.; Paton, M.B.; Reichardt, W.D.; Chu, K.C.; Linhart, M.S. (1980) Neoplastic and nonneoplastic lesions in aging Osborne-Mendel rats. *Toxicol Appl Pharmacol* 55:433-447.
- Graham, S.; Levin, M.L. (1971) Smoking withdrawal in the reduction of risk of lung cancer. *Cancer* 27:865-871.
- Greenberg, R.A.; Bauman, K.E.; Glover, L.H.; Strecher, V.J.; Kleinbaum, D.G.; Haley, N.J.; Steadman, H.C.; Fowler, M.G.; Loda, F.A. (1989) Ecology of passive smoking by young infants. *J Pediatr* 114:774-780.
- Grimmer, G.; Brune, H.; Dettbarn, G.; Naujack, K.W.; Mohr, U.; Wenzel-Hartung, R. (1988) Contribution of polycyclic aromatic compounds to the carcinogenicity of sidestream smoke of cigarettes evaluated by implantation into the lungs of rats. *Cancer Letters* 43:173-177.

06 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- Fischer, T.; Weber, A.; Grandjean, E. (1978) Air pollution due to tobacco smoke in restaurants. *Int Arch. Occup Environ Health* 41:267-280.
- Fleming, D.W.; Cochi, S.L.; Hightower, A.W.; Broome, C.V. (1987) Childhood upper respiratory tract infections: To what degree is incidence affected by day-care attendance? *Pediatrics* 79:55-60.
- Fong, P. (1982) The hazard of cigarette smoke to nonsmokers. *J. Biol. Phys.* 10:65.
- Freedman, D.A.; Navidi, W. (1987a) On the multistage model for carcinogenesis. Tech. Report. No. 47, University of California.
- Freedman, D.A.; Navidi, W. (1987b) On the risk of lung cancer for ex-smokers. Tech. Report. No. 135, University of California.
- Freedman, D.A.; Navidi, W.C. (1989) Multistage models for carcinogens. *Environ Health Perspect* 81:169-188.
- Fried, P.A.; Watkinson, B.; Dillon, R.F.; Dulberg, C.S. (1987) Neonatal neurological status in a low-risk population after prenatal exposure to cigarettes, marijuana, and alcohol. In: *J. Dev. Behav. Pediatr.* 8(6):318-326.
- Friedman, G.D.; Petitti, D.B.; Bawol, R.D. (1983) Prevalence and correlates of passive smoking. *Am J Public Health* 73:401.
- Galyskinova, V. (1964) 3-4 benzopyrene determination in the smoky atmosphere of social meeting rooms and restaurants: a contribution to the problem of the noxiousness of so-called passive smoking. *Neoplasma* 11:465-468.
- Gao, Y.; Blot, W.J.; Zheng, W.; Ershow, A.G.; Hsu, C.W.; Levin, L.I.; Zhang, R.; Fraumeni, J.F. (1987) Lung cancer among Chinese women. *Int'l J Cancer* 40:604-609.
- Gaffney, M.; Altshuler, B. (1988) Examination of the role of cigarette smoke in lung carcinogenesis using multistage models. *J. Nat'l Cancer Inst.* 80:925-31.
- Garfinkel, L. (1981) Time trends in lung cancer mortality among nonsmokers and a note on passive smoking. *J. Nat'l Cancer Inst.* 6:1061-1066.
- Garfinkel, L. (1984) Passive smoking and cancer: American experience. *Prev Med* 13:691-697.
- Garfinkel, L. (1985) Selection, follow-up, and analysis in the American Cancer Society Prospective Studies. Presented at a Workshop on the Selection, Follow-up, and Analysis in Prospective Studies held at the Waldorf-Astoria Hotel, New York, N.Y., October 3-5, 1983.
- Garfinkel, L.; Auerbach, O.; Joubert, L. (1985) Involuntary smoking and lung cancer: a case-control study. *J. Nat'l Cancer Inst.* 75:463-469.

4 05 NOV 90

DRAFT--DO NOT QUOTE OR CITE

- Grimmer, G.; Brune, H.; Deutsch-Wenzel, R.; Dettbarn, G.; Jacob, J.; Naujack, K.W.; Mohr, U.; Ernst, H. (1987) Contribution of polycyclic aromatic hydrocarbons and nitro-derivatives to the carcinogenic impact of diesel engine exhaust condensate evaluated by implantation into the lungs of rats. *Cancer Letters* 37:173-180.
- Hammond, E.C. (1966) Smoking in relation to the death rates of one million men and women. In: Haenszel, W., ed. *Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*. National Cancer Institute Monograph No. 19, Washington, D.C., pp. 127-204.
- Hammond, E.C.; Garfinkel, L. (1975) Aspirin and coronary heart disease: findings of a prospective study. *Br Med J* 2:269-271.
- Hammond, E.C.; Garfinkel, L.; Seidman, H.; Lew, E.A. (1976) 'Tar' and nicotine content of cigarette smoke in relation to death rates. *Environmental Research* 12:263-274.
- Harke, H. P. (1974) The problem of passive smoking. In: *The Influence of smoking on the CO concentration of office rooms*. *Int Arch Arbeitsmed* 33:199-206.
- Harlap, S.; Davies, A.M. (1974) Infant admissions to hospital and maternal smoking. *Lancet* 1(7857):529-533.
- Harmsen, H.; Effenberger, E. (1957) Tobacco smoke in public transportation, dwellings and work rooms. *Arch Hug* 141:383-400.
- Harley, N.H.; Pasternack, B.S. (1981) A model for predicting lung cancer risks induced by environmental levels of radon daughters. *Health Phys.* 40:307-316.
- Harley, N.H.; Pasternack, B.S. (1982) Environmental radon daughter alpha dose factors in five-lobed human lung. *Health Phys.* 42:788-799.
- Hasselbad, V.; Anderson, H.S.; Humble, C.G.; Graham, M.G. (1981) Indoor environmental determinants of lung function in children. *American Review of Respiratory Disease* 123(5):479-485.
- Henderson, F.W.; Reid, H.F.; Morris, R.; Wang, O.; Hu, P.C.; Helms, R.W.; Forehand, L.; et al. (1989) Home air nicotine levels and urinary cotinine excretion in preschool children. *American Review of Respiratory Disease* 140:197-201.
- Henderson, F.W.; Morris, R.; Reid, H.F.; Hu, P.C. (1987) Serum and urine cotinine as quantitative measures of passive tobacco smoke exposure in young children. University of North Carolina, Chapel Hill, N.C.
- Herrmann, D.R.; Olsen, K.M.; Hiller, F.C. (1989) Nicotine absorption following pulmonary installation. Presented at the American Thoracic Society Meeting, Cincinnati, Ohio, May 1989.

06 NOV 90

DRAFT--DO NOT QUOTE OR CITE

- Hiller, F.C.; McCusker, K.T.; Mazumder, M.D.; Wilson, J.D.; Bone, R.C. (1982) Deposition of sidestream cigarette smoke in the human respiratory tract. *Am Rev Res Dis* 125:406-408.
- Hinds, W.C.; First, M.W. (1975) Concentrations of nicotine and tobacco smoke in public places. *New Engl J Med* 292:844-845.
- Hinton, A.E. (1989) Surgery for otitis media with effusion in children and its relationship to parental smoking. *J Laryngol Otol* 103(6):559-61.
- Hirayama, T. (1977) Smoking and cancer: A prospective study on cancer epidemiology based on a census population in Japan. In: *Smoking and Health*, vol. II. Washington DC, U.S. Public Health Service, DHEW Publ. No. (NIH)77-1413, pp. 65-72.
- Hirayama, T. (1981a) Non-smoking wives of heavy smokers have a higher risk of lung cancer: A study from Japan. *Br Med J* 282: 183-185.
- Hirayama, T. (1981b) (Letter) *Br Med J* 283:916-917.
- Hirayama, T. (1983a) Passive smoking and lung cancer: consistency of association. *Lancet* 2:1425-1426.
- Hirayama, T. (1983b) Passive smoking and lung cancer. Presented at the 5th World Conference on Smoking and Health, Winnipeg, Canada.
- Hirayama, T. (1984) Cancer mortality in nonsmoking women with smoking husbands based on a large-scale cohort study in Japan. *Prev Med* 13:680-690.
- Hoffmann, D.; Wynder, E.L. (1986) Chemical constituents and bioactivity of tobacco smoke. Zavidze, D.G. and Peto, R., eds. *Tobacco: A Major International Health Hazard*. Lyon, France: IARC Scientific Pub. No. 74, pp. 145-165.
- Hoffmann, D.; Wynder E.L.; Hecht, S.S.; Brunnemann, K.D.; LaVoie, E.J.; Haley, N.J. (1987) Chemical carcinogens in tobacco. Bannasch, P., ed. In: *Cancer Risks: Strategies for Elimination*. Berlin: Springer-Verlag, pp. 101-113.
- Hoffman, D.; Brunnemann, K.D.; Haley, N.J. (1989) Absorption of smoke constituents by nonsmokers. In: *Reducing Workplace Exposures to Environmental Tobacco Smoke*. EPA-HHS Manual (in press).
- Hoffmann, D.; Hecht, S.S. (1989) Advances in tobacco carcinogenesis. In: *Springer Handbook of Experimental Pharmacology: Chemical Carcinogenesis and Mutagenesis* (in press).
- Hole, D.J.; Gillis, C.R.; Chopra, C.; Hawthorne, V.M. (1989) Passive smoking and cardiorespiratory health in a general population in the west of Scotland. *British Med. Journal* 299:423-427.
- Hollander, M.; Wolfe, D.A. (1973) *Nonparametric statistical methods*. New York: John Wiley and Sons.



05 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- Houthuijs, D.; Remijn, B.; Brunekreef, B.; de Koning, R. (1987) Exposure to nitrogen dioxide and tobacco smoke and respiratory health of children. In: Indoor Air '87: Proceedings of The 4th International Conference on Indoor Air Quality and Climate 1:463-467.
- Hulka, B.S. (1988) The health consequences of environmental tobacco smoke. Environ Technol Lett 9(6):531-538.
- Humble, C.G.; Samet, J.M.; Pathak, D.R. (1987) Marriage to a smoker and lung cancer risk. Am J Public Health 77:598-602.
- Husgafvel-Pursiainen, K.; Sorsa, M.; Moller, M.; Benestad, C. (1986) Genotoxicity and polynuclear aromatic hydrocarbon analysis of environmental tobacco smoke samples from restaurants. Mutagenesis 1:287-291.
- Ingebretson, B.J.; Sears, S.B. (October 2-5, 1985) Particle size distribution measurements of sidestream cigarette smoke. Presented at the 39th Tobacco Chemists' Research Conference, Montreal, Canada.
- Inoue, R.; Hirayama, T. (1988) Passive smoking and lung cancer in women. In: Smoking and Health. Elsevier Science Publishers, pp. 283-285.
- International Agency for Research on Cancer (IARC). (1987) Environmental carcinogens-methods of analysis and exposure measurement Volume 9: Passive Smoking (O'Neill, I.K.; Brunnemann, K.D.; Dodet, B.; and Hoffmann, D. eds.) Lyon: IARC Scientific Publications No. 81.
- International Commission on Radiological Protection, ICRP Task Group on Lung Dynamics. (1966) Deposition and retention models for internal dosimetry of the human respiratory tract. Health Phys. 12:173-207.
- Ishizu, Y.; Ohta, K.; Okada, T. (1980) The effect of moisture on the growth of cigarette smoke particles. Beitr Tabakforsch 10:16.
- Jacob P.; Benowitz, N.; Alexander, T. (1988) Recent studies of nicotine metabolism in humans. Pharmacology Biochemistry and Behavior 30:249-253.
- Janoff, A.; Pryor, W.A.; Bengali, B.H. (1987) Effects of tobacco smoke components on cellular and biochemical processes in the lung. Am Rev Resp Dis 36:1058-1064.
- Jarvis, M.D.; Turnstall-Pedoe, H.; Feyerabend, C.; Vesey, C.; Saloojee, Y. (1984) Biochemical markers of smoke absorption and self-reported exposure to passive smoking. J Epid Comm Health 38:335-339.
- Jarvis, M.J.; Russell, M.A.; Feyerabend, C. et al. (1985) Passive exposure to tobacco smoke: Saliva cotinine concentrations in a representative population sample of nonsmoking school children. Br. J. Med 291:927-929.

05 JULY 80

DRAFT--DO NOT QUOTE OR CITE

- Jarvis, M.J.; Russell, M.A.H.; Feyerabend, C.; Eiser, J.R.; Morgan, M.; Gammage, P.; Gray, E.M. (1985) Passive exposure to tobacco smoke: Saliva cotinine concentrations in a representative population sample of nonsmoking schoolchildren. *Br. Med. J.* 291:929.
- Jarvis, M.J.; McNeill, A.D.; Russell, M.A.H.; West, A.; Bryant, A.; Geyerabend, C. (1987) Passive smoking in adolescents: One-year stability of exposure in the home. *Lancet* 1:1324-1325.
- Jarvis, M.; Russell, M.A.; Benowitz, N.L.; Feyerabend, C. (1988) Elimination of cotinine from body fluids: implications for noninvasive measurement of tobacco smoke exposure. *Am J Public Health* 78:696-698.
- Jarvis, M.J. (1989) Application of biochemical intake markers to passive smoking measurement and risk estimation. *Mutation Research* 222:101-110.
- Just J.; Borkowska, M.; Maziarka, S. (1972) Air pollution by tobacco smoke in Warsaw coffee houses. *Rocz Pantstw Zak Hig* 23:129-135.
- Kabat, G.C.; Wynder, E.L. (1984) Lung cancer in nonsmokers. *Cancer* 53:1214-1221.
- Kahn, H.A. (1966) The Dorn study of smoking and mortality among U.S. veterans: Report on eight and one-half years of observation, In: Haenszel, W., ed. *Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*. National Cancer Institute Monograph No. 19, Washington, D.C., pg. 1.
- Kallail, K.J.; Rainbolt, H.R.; Bruntzel, M.D. (1987) Passive smoking and middle ear problems in Kansas public school children. *J. Commun. Disord.* 20:187-196.
- Kasuga, H.; Hasabe, A.; Matsuki, H.; Osaka, F. (1979) Respiratory symptoms in school children and the role of passive smoking. *Tokai Journal of Experimental and Clinical Medicine* 4(2):101-114.
- Kauffmann, F.; Oriol, P.; Tessier, J.-F. (1983) Adult passive smoking in the home environment: A risk factor for chronic airflow limitation. *American Journal of Epidemiology* 117(3):269-280.
- Kauffmann, F.; Tager, I.B.; Munoz, A.; Speizer, F.E. (1989) Familial factors related to lung function in children aged 6-10 years: Results from the PAARC epidemiologic study. *Am J Epidemiol* 129(6):1289-1299.
- Keith, C.H.; Derrick, J.D. (1960) Measurement of the particle size distribution and concentration of cigarette smoke by the confuge. *J Colloid Sci* 15:340-356.
- Kentner, M.; Triebig, G.; Weltle, D. (1984) The influence of passive smoking on pulmonary function: A study of 1351 office workers. *Preventive Medicine* 13(6):656-669.
- Kilpatrick, S.J. (1987) Misclassification of environmental tobacco smoke exposure: its potential influence on studies of environmental tobacco smoke and lung cancer. *Toxicology Letters* 35:163-168.

05 10/17 50

DRAFT--DO NOT QUOTE OR CITE

- Kleinman, J.C.; Pierre, M.B.; Madans, J.F.; Land, G.H.; Schramm, W.F. (1988) The effects of maternal smoking on fetal and infant mortality. *Am. J. Epidemiol.* 127(2):274-282.
- Knoth, A.; Bohn, H.; Schmidt, F. (1983) Passive smoking as a causal factor of bronchial carcinoma in female nonsmokers. *Medizinisch Klin* 78:66-69.
- Koo, L.C.; Ho, J.H.; Saw, D.; Ho, C.Y. (1987) Measurements of passive smoking and estimates of lung cancer risk among non-smoking Chinese females. *Int J Cancer* 39:162-169.
- Koo, L.C.; Ho, J.H.-C.; Matsuki, H.; Shimizu, H.; Mori, T.; Tominaga, S. (1988) A comparison of the prevalence of respiratory illnesses among nonsmoking mothers and their children in Japan and Hong Kong. *Am Rev Respir Dis* 138:290-295.
- Kraemer, M.J.; Bierman, C.W.; Furukawa, C.T.; Pierson, W.E.; Richardson, M.A.; Shapiro, G.G.; Weiss, N.S. (1983) Risk factors for persistent middle-ear effusions: Otitis media, catarrh, cigarette smoke exposure, and atopy. *Journal of the American Medical Association* 249(8):1022-1025.
- Kraus, J.F.; Greenland, S.; Bulterys, M. (1989) Risk factors for sudden infant death syndrome in the US Collaborative Perinatal Project. *Int. J. Epidemiol.* 18(1):113-120.
- Kuller, L.H.; Garfinkel, L.; Correa, P.; Haley, N.; Hoffmann, D.; Preston-Martin, S.; Sandler, D. (1986) Contribution of passive smoking to respiratory cancer. *Environ Health Perspectives* 70:57-69.
- Kyermater, G.A.; Damiano, M.D.; Dovorchik, B.H.; Vesell, D.S. (1982) Smoking-induced changes in nicotine disposition: application of a new HPLC assay for nicotine and its metabolites. *Clin-Pharm Ther* 32:769-780.
- Lam, T. H.; Kung, I.T.M.; Wong, C.M.; Lam, W.K.; Kleevens, J.W.L.; Saw, D.; Hsu, C.; Seneviratne, S.; Lam, S.Y.; Lo, K.K.; Chan, W.C. (1987) Smoking, passive smoking and histological types in lung cancer in Hong Kong Chinese women. *Br J Cancer* 6:673-378.
- Lam, W.K. (1985) A clinical and epidemiological study of carcinoma of lung in Hong Kong. Thesis submitted to the University of Hong Kong for the degree of Doctor of Medicine.
- Layard, M.W.; Viren, J.R. (1989) Assessing the validity of a Japanese cohort study. Elsevier Science Publishers B.V. (Biomedical Division).
- Leaderer, B.P. (1988) Measuring exposure to environmental tobacco smoke. Report prepared for the U. S. Environmental Protection Agency.
- Leaderer, B.P.; Cain, W.S.; Isseroff, R.; Berglund, L.G. (1984) Ventilation requirements in buildings II. Particulate matter and carbon monoxide from cigarette smoking. *Atmos. Environ.* 18, 99-106.
- Leaderer, B. (1990) Assessing exposure to environmental tobacco smoke. *Risk Analysis*, Vol. 10, No. 1.

06 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- Lebowitz, M.D.; Knudson, R.J.; Burrows, B. (1975) The Tucson epidemiology study of chronic obstructive lung disease. I. Methodology and prevalence of disease. *Am J. Epidemiol.* 102:137-152.
- Lebowitz, M.D.; Burrows, B. (1976) Respiratory symptoms related to smoking habits of family adults. *Chest* 69(1):48-50.
- Lebowitz, M.D.; Armet, D.B.; Knudson, R. (1982) The effect of passive smoking on pulmonary function in children. *Environment International* 8(1-6):371-373.
- Lebowitz, M.D.; Holberg, C.J.; Knudson, R.J.; Burrows, B. (1987) Longitudinal study of pulmonary function development in childhood, adolescence, and early adulthood. *Am. Rev. Resp. Dis.* 136:69-75.
- Lebowitz, M.D.; Holberg, C.J. (1987) Physiological changes in pulmonary development related to passive smoking and its interaction with active smoking. *Toxicology Letters* 35:101-105.
- Lebowitz, M.D.; Holberg, C.J. (1988) Effects of parental smoking and other risk factors on the development of pulmonary function in children and adolescents. *American Journal of Epidemiology* 128(3):589-597.
- Lee, B.L.; Benowitz, N.L.; Jacob P. (1987) Influence of tobacco abstinence of the disposition kinetics and effects of nicotine. *Clin Pharm Ther* 41:474-479.
- Lee, P.N.; Chamberlain, J.; Alderson, M.R. (1986) Relationship of passive smoking to risk of lung cancer and other smoking-associated diseases. *Br J Cancer* 54:97-105.
- Lee, P.N. (1987a) Lung cancer and passive smoking: Association an artefact due to misclassification of smoking habits. *Toxicology Letters* 35:157-162.
- Lee, P.N. (1987b) Passive smoking and lung cancer association: a result of bias? *Human Toxicol* 6:517-524.
- Lee, P.N. (1988) Misclassification of smoking habits and passive smoking. Berlin: Springer
- Leeder, S.R.; Corkhill, R.T.; Irwig, L.M.; Holland, W.W. (1976a) Influence of family factors on asthma and wheezing during the first five years of life. In: *British Journal of Preventive and Social Medicine* 30(4):213-218.
- Leeder, S.R.; Corkhill, R.T.; Irwig, L.M.; Holland, W.W. (1976b) Influence of family factors on the incidence of lower respiratory illness during the first year of life. *British Journal of Preventive and Social Medicine* 30(4):203-212.
- Lehnert, G. (1984) Roundtable discussion from Symposium "Medical Perspectives on Passive Smoking." *Prev Med* 13:732-733.
- Leung, H.M.; Kupper, L.L. (1981) Comparisons of confidence intervals for attributable risk. *Biometrics* 37:293-302.

4 05 NOV 90

DRAFT: DO NOT QUOTE OR CITE

- Lindblad, A.; Marsal, K.; Andersson, K.-E. (1988) Effect of nicotine on human fetal blood flow. *Obstet. Gynecol.* 72(3):371-382.
- Lubin, J.H., Blot, W.J., Berrino, F., Flamant, R., Gillis, C.R., Kunze, M., Schmahl, D., Visco, G. (1984) Modifying risk of developing lung cancer by changing habits of cigarette smoking. *Br Med J* 288:1953-1956.
- Luchsinger, P.C.; LaGarde, B.; Kilfeathers, J.E. (1968) Particle clearance from the human tracheobronchial tree. *Am. Rev Resp Dis* 97:1046.
- Mantel, N. (1963) Chi-square tests with one degree of freedom: extensions of the Mantel-Haenszel procedure. *J Am Statist Assoc* 58:690-700.
- Mantel, N. (1987a) Lung cancer and passive smoking. (Letter) *Br J Med* 294:440-441.
- Mantel, N. (1987b) Re: Passive smoking and lung cancer in Swedish women (Letter) *Am J Epidemiol* 126(6):1214-1216.
- Mantel, N.; Haenszel, W. (1959) Statistical aspects of the analysis of data from retrospective studies of disease. *J Natl Cancer Inst* 22:719-748.
- Maritz, G.S. (1988) Effect of maternal nicotine exposure on growth in vivo of lung tissue of neonatal rats. *Biol. Neonate, Foetal Neonatal Res.* 53(3):163-170.
- Marks, B.E. (1988) Respiratory illness and home environment. *British Medical Journal* 296(6638):1740.
- Martin, T.R.; Bracken, M.B. (1986) Association of low birth weight with passive smoke exposure in pregnancy. *American Journal of Epidemiology* 124(4):633-642.
- Martinez, F.D.; Antognoni, G.; Macri, F.; Bonci, E.; Midulla, F.; De Castro, G.; Ronchetti, R. (1988) Parental smoking enhances bronchial responsiveness in nine-year-old children. *Am. Rev. Respir. Dis.* 138(3):518-523.
- Martonen, T.; Hofmann, W. (1986) Factors to be considered in a dosimetry model for risk assessment of inhaled particles. *Radiation Protection Dosimetry* 15:225-232.
- Masi, M.A.; Hanley, J.A.; Ernst, P.; Becklake, M.R. (1988) Environmental exposure to tobacco smoke and lung function in young adults. *Am. Rev Resp Dis* 138:296-299.
- Matsuki, H.; Kasuga, H.; Osaka, F. et al. (1985) A comparative study on the health effects of smoking and indoor air pollution in summer and winter. In: *Tokai Journal of Experimental and Clinical Medicine* 10(4):427-437.
- Matsukura, S.; Taminato, T.; Kitano, N. et al. (1984) Effects of environmental tobacco smoke on urinary cotinine excretion in nonsmokers. *N Engl J Med* 311:828-832.
- McConnochie, K.M.; Roghmann, R.J. (1986) Breast feeding and maternal smoking as predictors of wheezing in children age 6 to 10 years. *Pediatr Pulmonol* 2:260-268.

05 NOV 90

DRAFT--DO NOT QUOTE OR CITE

McDowell, E.M.; Trump, B.F. (1983) Histogenesis of preneoplastic and neoplastic lesions in tracheobronchial epithelium. *Survey and Synthesis of Path. Res.*, 2, 235-279.

Medina, E.; Medina, R.; Kaempffer, A.M. (1988) Smoking and incidence of respiratory diseases in children. *Rev. Chil. Pediatr.*, 59(1):60-64.

Meyer, M.B.; Tonascia, J.A.; Buck, C. (1974) The interrelationship of maternal smoking and increased perinatal mortality with other risk factors. Further analysis of the Ontario perinatal mortality study, 1960-1961. *American Journal of Epidemiology* 100:443-452.

Miettinen, O.S. (1974) Proportion of disease caused or prevented by a given exposure, trait or intervention. *Am J Epidemiol* 99: 325-332.

Miller, G.H. (1984) Cancer, passive smoking and nonemployed and employed wives. *Western J Med* 140(4):632-635.

Mitchell, R.I. (1962) Controlled measurement of smoke particle retention in the respiratory tract. *Am Rev Resp Dis* 85:526-533.

Moghissi, A.A.; Seiler, F.A.; Dellarco, M.J. (1987) Enhancement of exposure to radon progeny as a consequence of passive smoking. *Proceedings of the 4th International Conference on Indoor Air Quality and Climate, West Berlin, Volume 2, August 1987*, pp. 360-364.

Moolgavkar, S.H.; Dewanji, A.; Luebeck, G. (1989) Cigarette smoking and lung cancer: reanalysis of the British doctors data. *J Am Cancer Inst.* 81(6):415-420.

Moorhead, R. (1985) Passive smoking and young children's health. *Australian Family Physician* 14(10):1058-1062.

Muramatsu, M.; Umemura, S.; Okada, T.; Tomita, H. (1984) Estimation of personal exposure to tobacco smoke with a newly developed nicotine personal monitor. *Environ Res* 35:218-227.

Murray, A.B.; Morrison, B.J. (1986) The effect of cigarette smoke from the mother on bronchial responsiveness and severity of symptoms in children with asthma. *J Allergy Clin Immunol* 77:575-581.

Murray, A.B.; Morrison, B.J. (1989) Passive smoking by asthmatics: Its greater effect on boys than on girls and on older than on younger children. *Pediatrics* 84:451-459.

Murray, A.B.; Morrison, B.J. (1988) Passive smoking and the seasonal difference of severity of asthma in children. *Chest* 94:701-708.

Muscatti, S.K.; Mackey, M.A.; Newsom, B. (1988) The influence of smoking and stress on prenatal weight gain and infant birth weight of teenage mothers. *J. Nutr. Educ.* 20(6):299-302.

Nash, J.E.; Persaud, T.V.N. (1988) Influence of nicotine and caffeine on rat embryonic development.

05 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- National Academy of Sciences. (1988) Health risks of radon and other internally deposited alpha-emitters, BEIR IV, National Academy Press, Washington, D.C.
- National Council on Radiation Protection and Measurement (NCRP) (1984) Evaluation of occupational and environmental exposures to radon daughters in the United States. NCRP Report No. 78, Bethesda, MD.
- National Research Council (NRC) (1986a) Environmental tobacco smoke: Measuring exposures and assessing health effects, National Academy Press, Washington, D.C.
- National Research Council (NRC) (1986b) Air quality and safety: The airliner cabin environment. National Academy Press, Washington, DC.
- Nazaroff, W.W.; Cass, G.R. (1989) Mathematical modeling of indoor aerosol dynamics. Environ Sci Technol 23:157-166.
- Neal, A.D.; Wadden, R.A.; Rosenberg, S. (1978) Evaluation of indoor particulate concentrations for an urban hospital. Am Ind Hyg Assoc J 29:578-582.
- Nelson, P.A.; Qvant, F.R.; Sem, G.F. (1982) Experimental measurements of aerosol concentrations in offices. Environ Int 8:223-227.
- Neuspiel, D.R.; Rush, D.; Butler, N.R.; Golding, J.; Bijur, P.E.; Kurzon M. (1989) Parental smoking and post-infancy wheezing in children: A prospective cohort study. Am J Public Health 79(2):168-171.
- O'Connor, G.; Weiss, S.T.; Tager, I.; Speizer, F.E. (1986) The effect of passive smoking on nonspecific bronchial responsiveness in a population sample of children and young adults. Clinical Research 34(2):581A (abstract).
- Ogston, S.A.; Florey, C.; Du V; Walker, C.H.M. (1987) Association of infant alimentary and respiratory illness with parental smoking and other environmental factors. Journal of Epidemiology and Community Health 41:21-25.
- Ostro, B.D. (1988) Estimating the risks of smoking, air pollution, and passive smoke on acute respiratory conditions. Risk Analysis 9(2):189-196.
- Paoletti, P.; Prediletto, R.; Carozzi, L.; Viegi, G.; Di Pede, F.; Carmignani, G.; Mammini, U.; Giuntini, C.; Lebowitz, M.D. (1989) Effects of childhood and adolescence-adulthood respiratory infections in a general population. Eur Respir J 2:428-436.
- Park, J.K.; Kim, I.S. (1986) Effects of family smoking on acute respiratory disease in children. Yonsei Med J 27(4):261-270.
- Pattishall, E.N.; Strobe, G.L.; Etzel, R.A.; Helms, R.W.; Haley, N.J.; Denny, F.W. (1985) Serum cotinine as a measure of tobacco smoke exposure in children. AJDC 139:1101-1104.

05 NOV 90

DRAFT--DO NOT QUOTE OR CITE

- Pedreira, F.A.; Guandolo, V.L.; Feroli, E.J.; Mella, G.W.; Weiss, I.P. (1985) Involuntary smoking and incidence of respiratory illness during the first year of life. *Pediatrics* 75(3):594-597.
- Pelech, L. (1989) Cooperation between the German Democratic Republic and the Czechoslovak socialist Republic in the study of long-term air pollution effects in selected groups of school-age children. *Z. Gesamte Hyg. Ihre Grenzgeb.* 35(1):50.
- Perry, J. (1973) Fasten your seat belts: no smoking. *Br Columbia Med J* 15:304-305.
- Pershagen, G.; Hrubec, Z.; Svensson, C. (1987) Passive smoking and lung cancer in Swedish women. *Am J Epidemiol* 125(1):17-24.
- Phillips, R.L.; Garfinkel, L.; Kuzma, J.W. et al. (1980a) Mortality among California Seventh-Day Adventists for selected cancer sites. *J Natl Cancer Inst* 65:1097-1107.
- Phillips, R.L.; Kuzma, J.W.; Beeson, W.L. et al. (1980b) Influence of selection versus lifestyle on risk of fatal cancer and cardiovascular disease among Seventh-Day Adventists. *Am J Epidemiol* 112: 296-314.
- Pimm, P.E.; Silverman, F.; Shephard, R.J. (1978) Physiological effects of acute passive exposure to cigarette smoke. *Archives of Environmental Health* 33(4):201-213.
- Plant, M.L.; Plant, M.A. (1988) Maternal use of alcohol and other drugs during pregnancy and birth abnormalities: Further results from a prospective study. *Alcohol Alcohol.* 23(3):229-233.
- Polydorva, M. (1961) An attempt to determine the retention of tobacco smoke by means of membrane filters. In: Davies, C.N. ed. *Inhaled Particles and Vapours* (Davies, C. N., ed.) Oxford: Pergamon Press, pp. 142-144.
- Portheine, F. (1971) To the problem of passive smoking. *Munch Med Wochenschr* 18:707-709.
- Pritchard, J.N.; Black, A.; McAughey, J.S. (1988) Physical behavior of sidestream smoke under ambient conditions. *Arch Environ Letters* 9(6): 545-552.
- Pullan, C.R.; Hey, E.N. (1982) Wheezing, asthma, and pulmonary dysfunction 10 years after infection with respiratory syncytial virus in infancy. *British Medical Journal* 284(6330):1665-1669.
- Pulkander, J.; Luotonen, J.; Timonen, M.; Karma, P. (1985) Risk factors affecting the occurrence of acute otitis media among 2-3 year-old urban children. *Acta Otolaryngologica* 100(3-4):260-265.
- Raabe, G.G. (1969) Concerning the interactions that occur between radon decay products and aerosols. *Health Phys.* 17, 177-187.
- Radsel, Z.; Kambic, V. (1978) The influence of cigarette smoke on the pharyngeal mucosa. *Acta. Otolaryngol.* 85:128-134.



41 06 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- Randerath, E.; Avitts, T.A.; Reddy, M.V.; Miller, R.H.; Everson, R.B.; Randerath, K. (1986) Comparative <sup>32</sup>P-analysis of cigarette smoke-induced DNA damage in human tissues and mouse skin. *Cancer Res* 46:5869-5877.
- Rantakallio, P. (1978) Relationship of maternal smoking to morbidity and mortality of the child up to the age of five. *Acta Paediatrica Scandinavica* 67(5):621-631.
- Redmond, C.K.; Ciocco, A.; Lloyd, J.W.; Rush, H.W. (1972) Long term mortality study of steelworkers. VI. Mortality from malignant neoplasms among coke oven workers. *J Occup Med* 14:621-629.
- Reed, B.D.; Lutz, L.J. (1988) Household smoking exposure--association with middle ear effusions. *Fam Med* 20(6):426-430.
- Remmer, H. (1987) Passively inhaled tobacco smoke: a challenge to toxicology and preventive medicine. *Arch Toxicol* 61:89-104.
- Repace, J.L. (1987) Indoor concentrations of environmental tobacco smoke: field surveys. In: O'Neill, I.K.; Brunemann, K.D.; Dodet B.; Hoffmann, D. eds. *Environmental Carcinogens: Methods of Analysis and Exposure Measurements Vol.9: Passive Smoking*. Lyon, France: IARC Scientific Pub. No. 81, pp. 141-162.
- Repace, J.L.; Lowrey, A.H. (1980) Indoor air pollution, tobacco smoke and public health. *Science* 208:464-472.
- Repace, J. L.; Lowrey, A. H. (1982) Tobacco smoke, ventilation and indoor air quality. *Am Soc Heat Refrig Air Cond Eng Trans* 88:894-914.
- Repace, J.L.; Lowrey, A.H. (1985) A quantitative estimate of nonsmokers' lung cancer risk from passive smoking. *Environ Int'l* 11:3-22.
- Repace, J.L.; Lowrey, A.H. (1987) Predicting the lung cancer risk of domestic passive smoking (Letter). *Amer Rev Resp Dis* 136:1308.
- Repace, J.L. (1989) Workplace restrictions on passive smoking: Justification on the basis of cancer risk (in press).
- Repace, J.L.; Lowrey, A.H. (1990) Risk assessment methodologies in passive smoking. *Journal of Risk Analysis* Vol. 10, No. 1.
- Riboli, E. (1987) Passive smoking and lung cancer: Epidemiological evidence and ongoing international collaborative studies. *Toxicology Letters* 35:19-27.
- Rickert, W.S.; Robinson, J.C.; Collinshaw, N.E. (1984) Sidestream yields of tar, nicotine and carbon monoxide from 15 brands of Canadian cigarettes. *Am J Pub Health* 74:228-231.
- Robins, J.M.; Blevins, D.; Schneiderman, M. (1989) The effective number of cigarettes inhaled daily by passive smokers: Are epidemiologic and dosimetric estimates consistent? *Journal of Hazardous Materials* 21:215-238.

7 06 NOV 90 CS

DRAFT--DO NOT QUOTE OR CITE

- Rogot, E. (1974) Smoking and mortality among U.S. veterans. *J Chronic Dis* 27:189-203.
- Rona, R.J.; Chinn, S.; Du Ve Florey, C. (1985) Exposure to cigarette smoking and children's growth. In: *International Journal of Epidemiology* 14(3):402-408.
- Rose, K.J.; Strand, F.L. (1987) Response of the developing neuromuscular system of the rat to nicotine and the neurotropic peptide fragment ACTH/MSH 4 - 10. *Ann. New York Acad. Sci.* 494:319-322.
- Rubin, B.K. (1986) Summer camp and passive smoking: Effects on nutrition, lung function and self-image in children with cystic fibrosis. *Chest* 89(6):480s.
- Rubin, D.H.; Damus, K. (1988) The relationship between passive smoking and child health: Methodologic criteria applied to prior studies. *Yale J Biol Med* 61(5):401-411.
- Rubin, D.H.; Krasilnikoff, P.A.; Leventhal, J.M.; Weile, B.; Berget, A. (1986) Effect of passive smoking on birth-weight. *The Lancet* Aug. 23.
- Rudnick, S.N.; Maher, E.F., (1986) Surface deposition of Rn-222 decay products with and without enhanced air motion. *Health Phys.* 51:283-293.
- Rush, D.; Cassano, P. (1970) Relationship of cigarette smoking and social class to birth weight and perinatal mortality among all births in Britain. *Journal of Epidemiology and Community Health* 37:249-55.
- Rush, D.; Cassano, P.; Harlap, S. (1988) Perinatal outcome, maternal weight gain, cigarette smoking and social status in Jerusalem. *Rev. Epidemiol. Sante Publ.* 36(3):186-195.
- Rush, D.; Callahan, K.R. (1989) Exposure to passive cigarette smoking and child development. A critical review. *Ann N Y Acad Sci* 562:74-100.
- Russell, M.A.H.; Jarvis, M.J.; West, R.J. (1986) Use of urinary nicotine concentrations to estimate exposure and mortality from passive smoking in non-smokers. *Br. J. Addiction* 81: 275-281.
- Rylander, B.; Eriksson, M.; Freyschuss, U. (1988) Risk factors from occasional and recurrent wheezing after RSV infection in infancy. *Acta Paediatr. Scand.* 77(5):711-715.
- Sackett, D.L. (1979) Bias in analytic research. *J Chronic Dis* 32(1-2):51-63.
- Sajd, G.; Lellouch, J.; Patois, E.; Zalokar, J. (1978) Parental smoking related to adenoidectomy and tonsillectomy in children. *Journal of Epidemiology and Community Health* 32(2):97-101.
- Samet, J.M.; Tager, I.B.; Speizer, F.E. (1983) The relationship between respiratory illness in childhood and chronic air-flow obstruction in adulthood. *American Review of Respiratory Disease* 127(4):508-523.

46 05 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- Samet, J.M. (1988a) Involuntary exposure to tobacco smoke. *Annals of Sports Medicine* 4(1):1-15.
- Samet, J.M. (1988b) Environmental tobacco smoke and cancer. Report prepared for the U.S. Environmental Protection Agency.
- Sandler, D.P.; Everson, R.B.; Wilcox, A.J. (1985) Passive smoking in adulthood and cancer risk. *Am J Epidemiol* 121(1):37-48.
- Saracci, R. (1989) Passive smoking and cancer risk. IARC report of panel of experts. Prepared at the request of the European School of Oncology through the Europe Against Cancer program of the European Economic Community.
- Saracci, R.; Riboli, E. (1989) Passive smoking and lung cancer: Current evidence and ongoing studies at the International Agency for Research on Cancer. *Mutation Research* 222:117-127.
- Schenker, M.B.; Samet, J.M.; Speizer, F.E. (1983) Risk factors for childhood respiratory disease: The effect of host factors and home environmental exposures. *American Review of Respiratory Disease* 128(6):1038-1043.
- Schesselman, J.J. (1982) Case-control studies: design, conduct, analysis. Monographs in Epidemiology and Biostatistics. New York: Oxford University Press, p. 190.
- Schilling, R.S.F.; Letai, A.D.; Hui, S.L.; Beck, G.J.; Schoenberg, J.B.; Bouhuys, A.H. (1977) Lung function, respiratory disease, and smoking in families. *American Journal of Epidemiology* 106(4):274-283.
- Schwartz-Bickenbach, D.; Schulte-Hobeln, B.; Abt, S.; Plum, C.; Nau, H. (1987) Smoking and passive smoking during pregnancy and early infancy: Effects on birth weight, lactation period, and cotinine concentrations in mother's milk and infant's urine. *Toxicology Letters* 35:73-81.
- Sebben, J.; Pimm, P.; Shephard, R. J. (1977) Cigarette smoke in enclosed public facilities. *Arch Envir Health* 32:53-58.
- Sepkrović, D.W.; Haley, N.J.; Hoffman, D. (1986) Elimination from the body of tobacco products by smokers and passive smokers. *J Am Med Assoc* 256:863.
- Shamsuddin, A.K.M.; Sinopoli, N.T.; Hemminki, K.; Boesch, R.R.; Harris, C.C. (1985) Detection of benzo[a]pyrene: DNA adducts in human white blood cells. *Cancer Res* 45:66-68.
- Shimizu, H.; Morishita, M.; Mizuno, K.; Masuda, T.; Ogura, Y.; Santo, M.; Nishimura, M.; Kunishima, K.; Karasawa, K.; Nishiwaki, K.; Yamamoto, M.; Hisamichi, S.; Tominaga, S. A case-control study of lung cancer in nonsmoking women. (1988) *Toboku J Exp Med* 154:389-397.

05 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- Sims, D.G.; Downham, M.A.P.S.; Gardner, P.S.; Webb, J.K.G.; Weightman, D. (1978) Study of 8-year-old children with a history of respiratory syncytial virus bronchiolitis in infancy. *British Medical Journal* 1(6104):11-14.
- Slavin, R.G.; Hertz M. (1975) Indoor air pollution: a study of the 30th annual meeting of the American Academy of Allergy. *Proceedings of the 30th Annual Academy of Allergy Conference*, p. 4.
- Somerville, S.M.; Rona, R.J.; Chinn, S. (1988) Passive smoking and respiratory conditions in primary school children. *Journal of Epidemiology And Community Health* 42(2):105-110.
- Speizer, F.E.; Ferris, B.Jr. Bishop, Y.M.M.; Spengler, J. (1980) Respiratory disease rates and pulmonary function in children associated with NO<sub>2</sub> exposure. *American Review of Respiratory Disease* 121(1):3-10.
- Speizer, F.E. (1989) Diagnosis of chronic obstructive pulmonary disease. *Am Rev of Respir Dis* 140:S106-S107.
- Spengler, J.D.; Dockery, D.W.; Turner, W.A.; Wolfson, J.M.; Ferris, B.G. (1981) Long-term measurements of respirable sulphates and particles inside and outside homes. *Atmos Environ* 15:23-30.
- Spinacci, S.; Arossa, W.; Burgiani, M.; Nortale, P.; Bucca, C.; DeCondisson, E. (1985) The effects of air pollution on the respiratory health of children: A cross-sectional study. In: *Pediatric Pulmonology* 1(5):262-266.
- Stanton, M.F.; Miller, E.; Wrench, C.; Blackwell, R. (1972) Experimental induction of epidermoid carcinoma in the lungs of rats by cigarette smoke condensate. *J Natl Cancer Inst* 49:867-877.
- Stehlik, G.; Rihter, O.; Altmann, H. (1982) Concentration of dimethylnitrosoamine in the air of smoke-filled rooms. *Ecotoxicol Environ Safety* 6:495-500.
- Sterling, T.D. (1983) Possible effects on occupational lung cancer from smoking related changes in the mucus content of the lung. *J Chronic Dis* 36:669-676.
- Stern, B.; Raizenne, M.; Burnett, R.; Kearney, J. (1987) Respiratory effects of early childhood exposure to passive smoke. *Indoor Air '87 Proceedings of the 4th International Conference on Indoor Air Quality and Climates*, Springer-Verlag New York, Inc.
- Stevens, R.G. and Moolgavkar, S.H. (1984) A cohort analysis of lung cancer and smoking in British males. *Am J Epidemiol* 119:624-641.
- Stober, W. (1984) Lung dynamics and uptake of smoke constituents by nonsmokers: a survey. *Prev Med* 13:589-601.
- Strachan, D.P.; Jarvis, M.J.; Feyerabend, C. (1989) Passive smoking, salivary cotinine concentrations, and middle ear effusion in 7 year old children. *British Medical Journal* 298:1549-1552.

4 05 NOV 80

DRAFT-DO NOT QUOTE OR CITE

- Svensson, C.; Pershagen, G.; Klominek, J. (1988) Smoking and passive smoking in relation to lung cancer in women. Department of Epidemiology, National Institute of Environmental Medicine, Stockholm, Sweden.
- Swift, D.L. (1989) Seminar presented at the University of North Carolina at Chapel Hill (unpublished).
- Szadkosi, D.; Harko, H.P.; Angerer, J. (1976) Burden of carbon monoxide from passive smoking in offices. *Int Med* 3:310-313.
- Tachi, N.; Aoyama, M. (1989) Effects of cigarette smoke exposure on early stage embryos in the rat. *Bull. Environ. Contam. Toxicol.* 43(3):467-472.
- Tager, I.B.; Munoz, A.; Rosner, B.; Weiss, S.T.; Carey, V.; Speizer, F.E. (1985) Effect of cigarette smoking on the pulmonary function of children and adolescents. *American Review of Respiratory Disease* 131(5):752-759.
- Tager, I.B.; Weiss, S.T.; Munoz, A.; Rosner, B.; Speizer, F.E. (1983) Longitudinal study of the effects of maternal smoking on pulmonary function in children. *New England Journal of Medicine* 309(42):699-703.
- Tager, I.B.; Weiss, S.T.; Rosner, B.; Speizer, F.E. (1979) Effect of parental cigarette smoking on the pulmonary function of children. *American Journal of Epidemiology* 110(1):15-26.
- Tager, I.B.; Segal, M.R.; Munoz, A.; Weiss, S.T.; Speizer, F.E. (1987) The effect of maternal cigarette smoking on the Pulmonary function of children and adolescents. *Am Rev Respir Dis* 136:1366-1370.
- Tager, I.B. (1986) Passive smoking and respiratory health in children: Sophistry or cause for concern? *American Review of Respiratory Disease* 133(6):959-961.
- Tainio, V.M.; Savilahti, E.; Salmenpera, L.; Arjomaa, P. (1988) Risk factors for infantile recurrent otitis media: Atopy but not type of feeding. *Pediatr. Res.* 23(5):509-512.
- Tashkin, D.; Clark, V.A.; Simmons, M.; Reems, C.; Coulson, A.H.; Bourque, L.B.; Sayre, J.W.; Detels, R.; Rokaw, S. (1984) The UCLA population studies of chronic obstructive respiratory disease: 7. Relationship between parental smoking and children's lung function. *American Review of Respiratory Disease* 129(6):891-897.
- Teculescu, D.B.; Pham, Q.T.; Varona-Lopez, W.; Deschamps J.P.; Marchand, M.; Henquel, J.C.; Manciaux, M. (1986) The single-breath nitrogen test does not detect functional impairment in children with passive exposure to tobacco smoke. *Bull. Eur. Physiopathol. Respir.* 22:605-607.
- Tenovu, A.H.; Kero, P.O.; Korvenranta, H.J.; Erkkola, R.U.; Klemi, P.J.; Tuominen, J. (1988) Risk factors associated with severely small for gestational age neonates. *Am. J. Perinatol.* 5(3):267-271.

4 06 NOV 90

DRAFT--DO NOT QUOTE OR CITE

- Thorslund, T.W.; Charnley, G. (1987) Use of the multistage model to predict the carcinogenic response associated with time-dependent exposure to multiple agents. Proceedings of the ASA/EPA Conference on Current Assessment of Combined Toxicants. EPA-230-03-87-027. Washington, D.C., Environmental Protection Agency, pp. 9-15.
- Thorslund, T.W. (1990) Task 115- Obtaining unit risk estimates for inhaled B[a]P. Task 116 - Development of a systematic approach for estimating PAH relative potencies. Prepared for the Office of Health and Environmental Assessment. Under EPA contract no. 68-02-4403. Preliminary Report.
- Thyssen, J.; Altoff, J.; Kimmerle, G.; Mohr, U. (1981) Inhalation studies with benzo[a]pyrene in Syrian golden hamsters. *J Natl Cancer Inst* 66:575-577.
- Tredaniel, J.; Hill, C.; Chastang C.; Hirsch A. (1989) Passive smoking and respiratory diseases. Current data. *Rev Mal Respir* 6(2):109-120.
- Trichopoulos, D. (1988) Passive smoking and lung cancer. *Scand J Soc Med* 16:75-79.
- Trichopoulos, D.; Kalandidi, A.; Sparros, L. (1983) Lung cancer and passive smoking: conclusion of Greek study. (Letter) *Lancet* 667-668.
- Tzoumaka-Bakoula, C.; Lekea-Karanika, V.; Matsaniotis, N.S.; Goldin, J. (1989) The Greek national perinatal survey. II: Socioeconomic factors and perinatal mortality in Greece. *Paediatr. Perinat. Epidemiol* 3(1):41-52.
- Überla, K. (1987) Lung cancer from passive smoking: hypothesis or convincing evidence? *Int Arch Envir Occup Health* 59:421-437.
- U.S. Department of Health and Human Services. (1984) The health consequences of smoking: Chronic obstructive lung disease. A Report of the Surgeon General. DHHS Pub. No. (PHS)84-50205. U.S. Department of Health and Human Services, Public Health Service, Office of the Assistant Secretary for Health, Office on Smoking and Health, 1984.
- U.S. Department of Transportation. (1971) Health aspects of smoking in transport aircraft. U.S. DOT, Washington, DC, National Technical Information Service.
- U.S. Environmental Protection Agency (EPA). (1984) Summary of the issues with regard to the carcinogenicity, mutagenicity, and exposure of polycyclic organic matter. NTIS No. PB88-220371.
- U.S. Surgeon General (U.S. SG) (1986) The health consequences of involuntary smoking. U.S. Department of Health and Human Services, Public Health Service.
- U. S. Surgeon General (U.S. SG). (1988) The health consequences of smoking: nicotine addiction. U.S. Department of Health and Human Services, Public Health Service.
- U.S. Surgeon General (U.S. SG) (1989) Reducing the health consequences of smoking. U.S. Department of Health and Human Services, Public Health Service.

06 NOV 80

DRAFT--DO NOT QUOTE OR CITE

- Vainio, H.; Partanen, T. (1989) Population burden of lung cancer to environmental tobacco smoke. *Mutation Research* 222:137-140.
- Vandenbroucke, J.P. (1988) Passive smoking and lung cancer: a publication bias? *Br J Med* 296:391-392.
- Varela, L.R. (1987) Assessment of the association between passive smoking and lung cancer. Dissertation to Yale University in candidacy for the degree of Doctor of Philosophy.
- Vedal, S.; Schenker, M.B.; Samet, J.M.; Speizer, F.E. (1984) Risk factors for childhood respiratory disease: Analysis of pulmonary function. *American Review of Respiratory Disease* 130(2):187-192.
- Vosmar, M.B.J.G.; Jongsma, H.W.; Van Dongen, P.W.J. (1989) The value of ultrasonic placental grading: No correlation with intrauterine growth retardation or with maternal smoking. *J. Perinat. Med.* 17(2):137-143.
- Vutuc, C. (1984) Quantitative aspects of passive smoking and lung cancer. *Prev Med* 13: 698-704.
- Wald, N.J.; Idle, M.; Boreham, J.; Bailey, A.; Van Vunatcis, H. (1981) Serum cotinine levels in pipe smokers: evidence against nicotine as a cause of coronary heart disease. *Lancet*, 2:775-777.
- Wald, N.J.; Nanchahal, K.; Thompson, S.G.; Cuckle, H.S. (1986) Does breathing other people's tobacco smoke cause lung cancer? *Br J Med* 293:1217-1222.
- Wald, N.J.; Ritchie, C. (1984) Validation of studies on lung cancer in nonsmokers married to smokers. *Lancet* 1:1067.
- Wald, N.; Doll, R.; Darby, S.; Kiryluk, S.; Peto, R.; Pike, M. eds. (1988) *U.K. Smoking Statistics*. Oxford University Press.
- Walter, S.D. (1975) The distribution of Levin's measure of attributable risk. *Biometrika* 62: 371-374.
- Walter, S.D. (1976) The estimation and interpretation of attributable risk in health research. *Biometrics* 32:829-849.
- Wanner, W.A.; Hirsch, J.A.; Greenelch, D.E.; Swenson, E.W.; Fore T. (1973) Tracheal mucus velocity in beagles after chronic exposure to cigarette smoke. *Arch Environ Health* 27:370-371.
- Ware, J.H.; Dockery, D.W.; Spiro, A.III; Speizer, F.E.; Ferris, B.G. Jr (1984) Passive smoking, gas cooking, and respiratory health of children living in six cities. *American Review of Respiratory Disease* 129(3) 366-374.
- Weber, A.; Fischer, T. (1980) Passive smoking at work. *Int Arch Occup Environ Health* 47:209-221.

4. 06 NOV 50

DRAFT--DO NOT QUOTE OR CITE

- Weber A.; Fischer T.; Grandjean, A. (1979) Passive smoking in experimental and field conditions. *Environ Resp.* 20:205-216.
- Weibel, E.R. (1963) Morphometry of the human lung. New York, NY: Academic Press.
- Weiss, S.T.; Tager, I.B.; Speizer, F.E.; Rosner, B. (1980) Persistent wheeze: Its relation to respiratory illness, cigarette smoking, and level of pulmonary function in a population sample of children. *American Review of Respiratory Disease* 122(5):697-707.
- Weiss, S.T.; Tager, I.B.; Schenker, M.; Speizer, F.E. (1983) The health effects of involuntary smoking. *Am Rev Respir Dis* 128:933-942.
- Weiss, T.; Dorow, P.; Felix, R. (1983) Regional mucociliary removal of inhaled particles in smokers with small airway disease. *Respiration* 44:338-345.
- Weiss, S.T. (1986) Passive smoking and lung cancer: what is the risk? (Editorial) *Am Rev Respir Dis* 33:1-3.
- Weitzman, M.; Gortmaker, S.; Walker, D.K.; Sobol, A. (1990) Maternal smoking and childhood asthma. *Pediatrics*, April 1990, 505-511.
- Wells, A.J. (1988a) Re: Passive smoking and lung cancer: a publication bias? (Letter) *Br Med J* 296:1128.
- Wells, A.J. (1988) An estimate of adult mortality in the United States from passive smoking. *Environ. Int'l* 14:249-265.
- Wells, A.J. (1986). Misclassification as a factor in passive smoking risk. (Letter) *Lancet* 2:638
- Wells, A.J. (1990) An estimate of adult mortality in the United States from passive smoking: A response to criticism. *Environment International*, Vol. 16.
- White, J.R.; Froeh, H.F. (1980) Small-airways dysfunction in nonsmokers chronically exposed to tobacco smoke. *New England Journal of Medicine* 302(13):720-723.
- Whittemore, A.S. (1982) Statistical methods for estimating attributable risks from retrospective data. *Stat Med* 1:229-243.
- Whittemore, A.S. (1988) Effect of cigarette smoking in epidemiological studies of lung cancer. *Stat Med* 7:223-238.
- Wiedemann, H.P.; Mahler, D.A.; Loke, J.; Virguito, J.A.; Snyder, P.; Matthay, R.A. (1986) Acute effects of passive smoking on lung function and airway reactivity in asthmatic subjects. *Chest* 89(2):180-185.
- Wigle, D.T.; Collishaw, N.E.; Kirkbride, J.; Mao, Y. (1987) Deaths in Canada from lung cancer due to involuntary smoking. *J Can Med Assoc* 136: 945.



06 NOV 90

DRAFT--DO NOT QUOTE OR CITE

- Willatt, D.J. (1986) Children's sore throats related to parental smoking. *Clinical Otolaryngol* 11:417-321.
- Witorsch, R.I.; Witorsch, P. (1989) A critical analysis of the relationship between parental smoking and pulmonary performance in children. *Off Gesundheitswes* 51(2):78-83.
- Woolf, B. (1955) On estimating the relationship between blood group and disease. *Ann Human Genet* 19:251-253.
- World Health Organization (WHO). (1986) Principles for evaluating health risks from chemicals during infancy and early childhood: The need for a special approach. *Environmental Health Criteria* 59.
- Wu, A.H.; Henderson, B.E.; Pike, M.D.; Yu, M.C. (1985) Smoking and other risk factors for lung cancer in women. *J Natl Cancer Inst* 74(4):747-751.
- Wu, A.; Samet, J. Environmental tobacco smoke: Exposure-response relationships in epidemiologic studies. *Risk Analysis* Vol. 10, No. 1.
- Wynder, E.L.; Hoffman, D. (1967) Tobacco and tobacco smoke: studies in experimental carcinogenesis. New York: Academic Press.
- Yarnell, J.W.G.; St. Leger, A.S. (1979) Respiratory illness, maternal smoking habit and lung function in children. *British Journal of Diseases of the Chest* 73(3):230-236.
- Yusuf, S.; Petó, S.R.; Lewis, J.; Collins, R.; Sleight, P. (1985) Beta blockade during and after myocardial infarction: an overview of the randomized trials. *J Prog Cardiovasc Dis* 27:335-371.
- Zielhuis, G.A.; Heuvelmans-Heinen, E.W.; Rach, G.H.; Van den Broek, P. (1989) Environmental risk factors for otitis media with effusion in preschool children. *Scand. J. Prim. Health Care* 7(1):33-38.